

New data on Enidae (Gastropoda, Pulmonata) of Nepal

Andrey G. KUZNETSOV*, Anatoly A. SCHILEYKO**

*Sumskoj Proezd 2/3-63, Moscow, 113208, RUSSIA

**A. N. Severtzov Institute of Problems of Evolution, Russian Academy of Sciences, Leninski Prospect 33, Moscow 117071, RUSSIA

Pupinidius tukuchensis sp. nov., *Laevozebrinus nepalensis myagdiensis* subsp. nov., and *L. mustangensis* sp. nov. are described by conchological and anatomical features. New data on the distribution and habitat of *Nepaliena ceratina* (Benson, 1849), *Laevozebrinus nepalensis nepalensis* Schileyko et Frank, 1994, and *Mirus* (?) *nilagiricus* (Pfeiffer, 1846) are presented.

Новые данные о видах семейства Enidae фауны Непала

Андрей Г. КУЗНЕЦОВ*, Анатолий А. ШИЛЕЙКО**

*113208 Москва, Сумской проезд, 2/3-63

**Институт проблем экологии и эволюции им. А.Н.Северцова РАН, Москва 117071, Ленинский проспект, 33

Pupinidius tukuchensis sp. nov., *Laevozebrinus nepalensis myagdiensis* subsp. nov. и *L. mustangensis* sp. nov. описаны по конхологическим и анатомическим признакам. Для трех видов [*Nepaliena ceratina* (Benson, 1849), *Laevozebrinus nepalensis nepalensis* Schileyko et Frank, 1994 и *Mirus* (?) *nilagiricus* (Pfeiffer, 1846)] приведены новые подробные данные о распространении и характере местообитаний.

About 20 species of Enidae are known from Indian part of Himalayas [Gude, 1914], while only two species have been described from Nepalesian Himalayas [Schileyko, Frank, 1994]. This is connected with the fact that the malacofauna of Nepal is still poorly known. During two collecting trips to Nepal of one of the authors (A. K.) in 1995 and 1996, new data on the distribution of previously known species were obtained, and two new species and one new subspecies were found. One of them, *Pupinidius tukuchensis* sp. nov., is the first representative of the genus from southern slope of Himalayas. The others — *Laevozebrinus nepalensis myagdiensis* subsp. nov. and *L. mustangensis* sp. nov. — are the second and the third members of the genus from Nepal. *Mirus* (?) *nilagiricus* (L. Pfeiffer, 1846) is described by only conchological features, so its generic position is not quite clear.

Family Enidae Woodward, 1903

Subfamily Pseudonapaeinae
Schileyko, 1978

Genus *Pupinidius* Moellendorff, 1901

Moellendorff, 1901: 341 (*Buliminus* subgen.; type species *Buliminus pupinidius* Moellendorff, 1901; o.d.); Wiegmann, 1901: 254; Schileyko, 1978: 345.

Pupinidius tukuchensis
Kuznetsov et Schileyko, sp. nov.

(Figs. 1, 2)

LOCUS TYPICUS — Western Nepal, Dhaulagiri zone, Mustang District, Annapurna National Park, Pholong-Dara Ridge, right side of Kali-Gandaki Valley, 200-700 m NE of Khobang village, 1-5 m over "Khobang —